

Table 3. Results for the amount and localization of TILs in Luminal B subtype of BC (n = 25)

Intratumoral	n=25 (%)	Stromal	n=25 (%)	χ^2	df	p
CD3				18.79	2	<0.0001
0	2 (8.0)					
Low	20 (80.0)	Low	7 (28.0)			
High	3 (12.0)	High	18 (72.0)			
CD4				12.52	2	0.0019
0	13 (52.0)		2 (8.0)			
Low	12 (48.0)	Low	21 (84.0)			
High		High	2 (8.0)			
CD8				10.48	2	0.0053
0	6 (24.0)		0 (0)			
Low	17 (68.0)	Low	16 (64.0)			
High	2 (8.0)	High	9 (36.0)			
FoxP3				5.69	2	0.0581
0	3 (12.0)	0	0 (0)			
Low	21 (84.0)	Low	20 (80.0)			
High	1 (4.0)	High	5 (20.0)			
CD20				14.19	2	0.0008
0	12 (48.0)	0	1 (4.0)			
Low	13 (52.0)	Low	21 (84.0)			
High	0 (0)	High	3 (12.0)			
TIL (%)						
≤ 50%			24 (96.0)			
> 50%			1 (4.0)			

Table 4. Results for the amount and localization of TILs in HER2-positive subtype of BC (n = 25)

Intratumoral	n=25 (%)	Stromal	n=25 (%)	χ^2	df	p
CD3				16.45	2	0.0003
0	1 (4.0)		1 (4.0)			
Low	20 (80.0)	Low	6 (24.0)			
High	4 (16.0)	High	18 (72.0)			
CD4				21.17	2	<0.0001
0	19 (76.0)		3 (12.0)			
Low	6 (24.0)	Low	20 (80.0)			
High	0 (0)	High	2 (8.0)			
CD8				10.29	2	0.0058
0	3 (12.0)		0 (0)			
Low	20 (80.0)	Low	14 (56.0)			
High	2 (8.0)	High	11(44.0)			
FoxP3				9.52	2	0.0086
0	4 (16.0)	0	2 (8.0)			
Low	20 (80.0)	Low	13 (52.0)			
High	1 (4.0)	High	10 (40.0)			
CD20				10.10	2	0.0064
0	6 (24.0)	0	0 (0)			
Low	19 (76.0)	Low	21 (84.0)			
High	0 (0)	High	4 (16.0)			
TIL (%)						
≤ 50%			20 (80.0)			
> 50%			5 (20.0)			

