

Original Article

STUDY OF THE CONTENT OF HEAVY METALS IN FOODS IN BULGARIA

Ivelina R. Drambozova

*Department of Hygiene, Medical Ecology,
Occupational Diseases and Disaster Medicine,
Faculty of Public Health,
Medical University - Pleven*

Corresponding Author:

Ivelina R. Drambozova
Department of Hygiene, Medical Ecology,
Occupational Diseases and Disaster Medicine,
Faculty of Public Health,
Medical University – Pleven
1, Saint Kliment Ohridski Str.
Pleven, 5800
e-mail: ivelina.ruseva@abv.bg

Received: October 14, 2021

Revision received: October 29, 2021

Accepted: November 25, 2021

Summary

Food safety is a priority for the European Union (EU). EU food safety policy covers all sectors of the food chain. In recent years, numerous studies have been conducted in Bulgaria on contaminants in plant and animal foods as one of the priority areas of food safety policy. Metals are part of the chemical pollutants in the environment. People are exposed to metals from the environment and food and water contaminated with metals. Researchers have found that the intake of metals with food is different and that food exposure is the primary way of metal exposure.

Keywords: heavy metals in food, food samples, food safety

Introduction

It is accepted that metal elements with a relative mass larger than 40 and a relatively high density (above 0.5 g / cm^3) are called heavy metals [1]. Some heavy metals (copper, zinc, iron, etc.) belong to the so-called trace elements because, in specific concentrations, they are vital for the functioning of the human body. These substances can have an adverse effect when their amount in the body exceeds what is needed. Other heavy metals (such as lead, cadmium, mercury, etc.) are not biologically necessary and have a detrimental effect on human health [2, 3]. Heavy metals can remain in the environment for a long time, as well as accumulate in organisms, which is a premise for their long-term toxic effects [4, 5, 6].

Heavy metals are chemical contaminants in the environment that occur in nature, both naturally and due to human activity. Heavy metal pollution affects not only the environment but also the organisms that inhabit it [7]. Consumption of food and drinking water is the main route for human exposure to heavy metals [8, 9, 10, 11].

Every year in Bulgaria, a national food control program for contaminants is planned and implemented, which is part of the multi-annual national control plan of the Republic of Bulgaria, prepared following the requirements of the legislation of the European Union [12].

Aim

The study aimed to analyze the levels of heavy metals cadmium (Cd), lead (Pb), tin (Sn), and mercury (Hg) in the food products offered on the Bulgarian market.

Methods and Materials

The study covered the period 2013-2019. We analyzed 2757 food samples for metal content (including cadmium, lead, tin, and mercury), as well as the trace elements copper and zinc. The samples were collected from 77 plant and animal foods, which had different contributions to the average daily intake of metals such as leafy vegetables, root crops, cereals, and legumes; animal foods - milk, dairy products, meat and offal used for food, fish and shellfish, special purpose foods. The samples were prepared for direct analysis in accredited laboratories.

The method of Graphite-furnace atomic absorption spectrometry (GF-AAS) was used to determine the metal content in the tested food samples (Figure 1).

The EU has strict food contamination standards to protect consumers' health [13, 14, 15, 16, 17, 18, 20]. Regulation (EC) № 1881/2006 (as amended by Regulation (EC) 2021/1323 of the Commission of August 10, 2021) and Ordinance № 5, of February 9, 2015, defined maximum levels (mg / kg) for heavy metals in foods [20, 21].

Results and Discussion

The analysis of the results showed that 97.71% of all tested food samples corresponded to the established maximum permissible amounts for heavy metals in food. [20, 21] Non-compliant samples accounted for 2.28% of all samples, and the discrepancy was only in terms of cadmium and lead content (Table 1).

From Table 1, it can be seen that in products of animal origin, in 97.63% of the samples, the levels of heavy metals corresponded to the statutory maximum levels. Non-compliant samples were only 2.36%.

Compared with the tested products of




Figure 1. Samples tested of foodstuffs for heavy metals content for the period 2013 – 2019

Table 1. Compliance of the tested samples with the established maximum permissible quantities for heavy metals in food

	Number of samples	Number of samples over LOQ	Number of samples with the value of the result following the maximum permissible quantities (mg/kg)	Number of samples with a result value above the maximum permissible quantities (mg/kg)
Samples of plant origin tested	1149	465	1124	25
Samples of animal origin tested	1608	1434	1570	38
Total	2757	1899	2697	60

vegetable origin, the percentage of non-standard samples in the products of animal origin was higher. In the tested samples of plant foods, 97.82% of them corresponded to the statutory maximum permissible amounts, while in 2.17%, it was found that these norms were exceeded (Figure 2).

Of the 77 types of food studied, 19.48% had a metal content above the maximum allowable amounts.

The simultaneous presence of several heavy metals in one food product was established. 6.49% of the studied types of food contain two metals - cadmium and lead, in a concentration above the maximum allowable amounts. It is established in apples, wild boar meat, herbs and spices, horse meat, and fresh spinach.

Only cadmium or lead was present in 12.98% of the tested foods, in concentrations above the permissible level.

In the studied types of food, the amount of mercury corresponded to the maximum regulated amount.

In the case of products of animal origin:

- 3.44% of the samples of wild boar meat;
- 5.76% of the cow's milk samples;
- 3.33% of the sheep's milk samples;
- 3.12% of the offal samples;
- 1.94% of honey samples;
- 26,31% of the samples of horsemeat;
- 11,11% of the samples of fish and other seafood did not meet the requirements of Regulation (EC) (1831/2003).

In all other tested products of animal origin, the quantities of heavy metals met the regulatory

requirements.

In the tested food products of plant origin, the share of non-standard samples was distributed as follows:

- 2.23% of the samples spinach canned/ deep-frozen;
- 2.72% of fresh spinach samples;
- 11.53% of the apple samples;
- 4.16% of the aubergine samples;
- 0.90% of carrot samples; 12.90% of the samples of herbs and spices;
- 5% of the samples Oyster mushroom sponge mushroom;
- 3.70% of the samples peppers, red.

Samples with established high concentrations of heavy metals are were single:

- lead in wild boar meat - 0.49 mg Pb / kg, standard 0.10 mg Pb / kg;

- lead in cow's milk - 0.1900 mg Pb / kg, standard 0.020 mg Pb / kg;

- lead in sheep's milk - 0.0970 mg Pb / kg, standard 0.020 mg Pb / kg;

- lead and cadmium in apples - 0.4900 mg Pb / kg, standard 0.10 mg Pb / kg and 0.4400 mg Cd / kg at a rate of 0.050 mg Cd / kg;

- cadmium in fish and other seafood - 0.1170 mg Cd / kg, standard 0.050 mg Cd / kg;

- cadmium in horse meat - 0.8990 mg Cd / kg, standard 0.20 mg Cd / kg;

- cadmium in spinach - 0.4500 mg Cd / kg, standard 0.20 mg Cd / kg.

Food was found to be one of the main sources of lead exposure. In Bulgaria, the consumption of bread and pasta is high. In children of all ages, the bread consumed is half the amount of cereals




Figure 2. Results for the content of heavy metals in the tested food samples




Figure 3. Results of compliance of the tested samples with the established maximum levels for heavy metals in food and samples with a value above the limit of quantification




Figure 4. Results of compliance of the tested samples with the established maximum levels for heavy metals in food and samples with a value above the limit of quantification

consumed, the average daily consumption of vegetables and milk is also slightly high and that of fish. Cereals, potatoes, leafy vegetables, milk, meat, offal, game consumption are the alimentary route of lead exposure [22].

The established amounts of tin (inorganic) in canned foods, other than beverages, canned fruits (peach and pineapple), fruit and vegetable juices, tomato paste were within the maximum allowable amounts.

In the studied fish samples, the amount of mercury corresponded to the determined maximum regulated amount of mercury in food.

The limit of quantification (LOQ) is the lowest amount of a substance that can be measured with certainty by standard laboratory tests [23, 24, 25, 26]. “The LOQ is calculated from the lowest concentration point taking into

account the recovery of internal standards and sample intake” [27].

The simultaneous presence of several heavy metals in the studied foods was found within the limits of quantification. Of all tested samples, 68.9% had a value above LOQ for more than two metals and trace elements.

For the samples of animal origin, 89.17% of the results were above LOQ, and for products of plant origin, 40.46% of the results were above LOQ for more than two metals and trace elements (Figure 3).

The most significant number of samples with a value above LOQ were found in samples of game meat, meat and offal of cattle, sheep, pigs and poultry, fish and bivalve molluscs, carrots, milk, honey, leafy vegetables (spinach), infusion of herbal tea, horse meat, pheasants and rabbit

meat.

All samples of sturgeon, beef kidney, veal liver, offal of farm animals, fish and other seafood, game birds, horse meat, mussels, mutton/lamb kidney, mutton/lamb liver, pork kidney, and rabbit meat were with values above LOQ. These products represented 20.77% of the tested foods and 8.56% of all samples.

With a value above LOQ were:

- 85.28% of the samples from wild boar,
- 95.34% from the samples from carp,
- 87.27% from the samples from carrots,
- 95.06% from the samples from chicken liver,
- 83.97% from the samples from cow's milk,
- 93.75% of the samples duck liver,
- 79.16 5 of the samples herbal tea infusion,
- 79.22% of the samples honey, 35.29% of the lettuce samples,
- 37.5% of the samples potatoes, 96.03% from pheasant meat samples,
- 97.18% from pork liver samples,
- 45% from rice samples, 96.05% from salmon and trout samples salmo spp.,
- 74.44% from sheep milk samples,
- 61.19% from spinach samples,
- 53.12% of the deer meat samples (Figure 4).

Discussion

It should be noted that children take more metals than adults, in the following order: Mn> Zn> Cu> Sr> Cr> Ni> As> Cd> Pb> Co> Sb. Lead is the most bioaccumulated metal of mussels and fish, along with cadmium. The sequence of accumulation of metals in different tissues of fish is established in the following order: liver> gills> gonads> muscles [28, 29, 30, 31, 32].

The study, covering six years, showed a generally low level of contamination of the studied foods with heavy metals.

Non-compliance with the maximum levels for heavy metals in foods for cadmium and lead alone was established.

The content of heavy metals in foods of animal origin was higher than in foods of plant origin.

More metals accumulate in wild boar meat, cow's milk, fish, and other seafood. It is important to note that frequent consumption of these foods can increase the amounts of heavy metals, which is important for public health.

In general, no significant contamination with heavy metals of the foods offered on the market in Bulgaria was established.

Conclusion

Chronic exposure to heavy metals through food intake can cause adverse health effects.

A risk assessment should be carried out so that actions and measures can be taken to reduce exposure levels and achieve safe exposure levels.

The results of the analysis can be used to assess the risk to human health associated with the detected content of heavy metals and serve in planning the monitoring and control of heavy metals in foods.

References

1. Bekyarov GL, Expert opinion Heavy metals - their impact on health, sources of pollution and non-clinical methods for their determination in the human body, 2009 October: 5-6.
2. WHO Assessment of chemical risks in food, <https://www.who.int/activities/assessing-chemical-risks-in-food/assessing-hemical-risks-in-food>
3. Stoyanov, S. Heavy metals in the environment and food products, toxic damage to humans, QC and prevention 1999; 954-14-0036-1:4-23.
4. European Guide to Good Hygiene Practices for the collection, storage, trading and transport of cereals, oilseeds, protein crops, other plant products and products derived thereof, Coceral/ Cogeca/Unistock Update Version 2.2 – July 2015:73.
5. Hartwig A, Jahnke GЛ Metals and their compounds as contaminants in food : Arsenic, cadmium, lead and aluminum, Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz, 2017; 60(7):715-21.
6. Suomi J, Tuominen P, Niinistö S, Virtanen S M and Savela Kp Dietary heavy metal exposure of Finnish 1-year-olds, AIMS Agriculture and Food, 4(3):778–93.
7. European Food Safety Authority (EFSA), Focalpoint Bulgaria, <https://focalpointbg.com>
8. Lopez-Alonso M, Rey-Crespo F, Herrero-Latorre C; Miranda M. Identifying sources of metal exposure in organic and conventional dairy farming, Chemosphere, 2017 Oct;185:1048-55.
9. Hu B, Deng F, Chen G, Chen X, Gao W, Long Le et al. Evolution of Absciscic Acid Signaling for Stress Responses to Toxic Metals and Metalloids, Front Plant Sci., 17 July 2020.

10. WHO Evaluation of certain food additives and contaminants the 53d report of the Joint FAO/WHO Expert Committee on Food Additives, 9789241209953, 2000:27-9.
11. WHO FOOD ADDITIVES SERIES: 63, FAO JECFA MONOGRAPHS 8, Safety evaluation of certain contaminants in food, 2011 FAO / WHO, 2006;EFSA, 2007a, b, 2008.
12. <https://www.mzh.government.bg/bg/politiki-i-programi/politiki-i-strategii/politiki-po-agrohranitelnata-veriga/edinennogogodishen-nacionalen-plan-za-kontrol/>
13. Codex Alimentarius Commission, General standard for contaminants and toxins in food and feed, CXS 193-1995, Adopted in 1995, Revised in 1997, 2006, 2008, 2009, Amended in 2010, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019.
14. Commission Regulation (EU) No 488/2014 of 12 May 2014 amending Regulation (EC) No 1881/2006 as regards maximum levels of cadmium in foodstuffs Text with EEA relevance.
15. Regulation (EC) No 178/2002 of the European Parliament and of the Council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety OJ L 31, 1.2.2002 r., p. 1–24.
16. Council Regulation (EEC) No 315/93 of 8 February 1993 laying down Community procedures for contaminants in food OJ L 37, 13.2.1993, p. 1–3.
17. Commission Regulation (EU) 2015/1005 of 25 June 2015 amending Regulation (EC) No 1881/2006 as regards maximum levels of lead in certain foodstuffs, OJ L 161, 26.6.2015, p. 9–13.
18. Commission Regulation (EU) 2021/1323 of 10 August 2021 amending Regulation (EC) No 1881/2006 as regards maximum levels of cadmium in certain foodstuffs, C/2021/5837, OJ L 288, 11.8.2021, p. 13–8.
19. Commission Regulation (EU) 2021/1323 of 10 August 2021 amending Regulation (EC) No 1881/2006 as regards maximum levels of cadmium in certain foodstuffs, C/2021/5837 OJ L 288, 11.8.2021, p. 13–8.
20. Commission Regulation (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs, OJ L 364, 20.12.2006, p. 5–24.
21. Ordinance № 5 of February 9, 2015 on determining the maximum permissible amounts of certain contaminants in food, effective from 20.02.2015 issued by the Minister of Health, prom. DV issue 14 of 20 February 2015, amended and ext. DV issue 11 of February 2, 2018.
22. National survey of nutrition factor for health risk among the population in Bulgaria, National survey of nutrition factor for health risk among the population in Bulgaria, Part III, 2014.
23. European Food Safety Authority (EFSA), Guidance Document on the Estimation of LOD and LOQ for Measurements in the Field of Contaminants in Feed and Food 2016.
24. European Food Safety Authority (EFSA) Scientific Committee, Guidance on selected default values to be used by the EFSA Scientific Committee, Scientific Panels and Units in the absence of actual measured data, EFSA Journal 2012;10(3):2579.
25. European Food Safety Authority (EFSA) The food classification and description system FoodEx2 (revision 2), 30 April 2015.
26. European Food Safety Authority (EFSA) Use of cut-off values on the limits of quantification reported in datasets used to estimate dietary exposure to chemical contaminants, 4 June 2018, doi:10.2903/sp.efsa.2018.EN-1452.
27. Commission Regulation (EU) No 589/2014 of 2 June 2014 laying down methods of sampling and analysis for the control of levels of dioxins, dioxin-like PCBs and non-dioxin-like PCBs in certain foodstuffs and repealing Regulation (EU) No 252/2012 Official Journal of the European Union 3.6.2014.
28. Milad Adel, Chiara Copat, Mohammad Reza Saeidi Asl, Gea Oliveri Conti, Mehdi Babazadeh, Margherita Ferrante, Bioaccumulation of trace metals in banded Persian bamboo shark from the Persian Gulf, 2018 Mar;113:198-203. doi: 10.1016/j.fct.2018.01.027. Epub 2018 Jan 31.
29. Figueira E. et al, Health concerns of consuming cockles from a low contaminated coastal system, July 2011, Environment International 37(5):965-72, DOI:10.1016/j.envint.2011.03.018, Source PubMed
30. Lopez-Alonso M. et al, Identifying sources of metal exposure in organic and conventional dairy farming, July 2017, Chemosphere 185.
31. Laird BD et al. Bioaccessibility of metals in fish, shellfish, wild game, and seaweed harvested in British Columbia 2013, Food and Chemical Toxicology, 58:381-7.
32. Wang R. et al. Concentrations and Exposure Evaluation of Metals in Diverse, 2018 Jan;74(1):131-9.