Journal of Biomedical and Clinical Research

Vol. 14, No. 2, Suppl. 1, 2021

EDITOR-IN-CHIEF

Grigor A. Gorchev Pleven, Bulgaria

DEPUTY EDITOR-IN-CHIEF

Mariya Sredkova Pleven, Bulgaria

EDITORIAL BOARD

Asparuh Asparuhov, Pleven, Bulgaria Chavdar Balabanov, Pleven, Bulgaria Margarita Alexandrova, Pleven, Bulgaria Tashko Deliyski, Pleven, Bulgaria Silviva Aleksandrova-Yankulovska, Pleven, Bulgaria Dimitar Gospodinov, Pleven, Bulgaria Emilyan Ivanov, Pleven, Bulgaria Yavor Ivanov, Pleven, Bulgaria Vaniva Nedkova, Pleven, Bulgaria Savelina Popovska, Pleven, Bulgaria Jordan Popov, Pleven, Bulgaria Dimitar Stoykov, Pleven, Bulgaria Mariya Simeonova, Pleven, Bulgaria Alexander Valkov, Pleven, Bulgaria Plamen Bojinov, Pleven, Bulgaria

EDITORIAL OFFICE

Assistant Editor Associated Editors

Statistical Editor Translation Editor Electronic Publishing Editor Production Editor

PUBLISHER

CO-EDITOR-IN-CHIEF

Slavcho T. Tomov Pleven, Bulgaria

ADVISORY BOARD

Genka Bekvarova, Varna, Bulgaria Veselin Borisov, Sofia, Bulgaria Yvon A. Carpentier, Bruxelles, Belgium Raimund Jakesz, Vienna, Austria Jorg Jerosh, Neuss, Germany Ivan Manchev, Stara Zagora, Bulgaria Pier P. Mariani, Rome, Italy Evlogi E. Marinov, Pleven, Bulgaria Robert P. Mecham, St. Louis, Missouri, USA Stewart Mennin, New Mexico, USA Ivan G. Mitov, Sofia, Bulgaria Gilbert Park, Cambridge, UK Penka Perenovska, Sofia, Bulgaria Peter M. Schlag, Berlin, Deutschland Emil Simeonov, Sofia, Bulgaria Svetla Torbova, Sofia, Bulgaria Nicola Vassilev, Sofia, Bulgaria Jacques Winand, Bruxelles, Belgium Hans P. Zenner, Tubingen, Germany

Radoslav Hristov Galya Stavreva Ivelina Yordanova Nadia Veleva Dobromir Dimitrov Petkana Hristova Yanka Tzvetanova Pencho Tonchev Silviya Emileva

Publishing center MU - Pleven

JBCR (ISSN 1313-6917) is a multidisciplinary peer-reviewed, open access scientific journal of Medical University-Pleven, published two times per year.

JBCR Online (ISSN 1313-9053) offers free access to all articles at jbcr.mu-pleven.bg **Editorial Office:** Journal of Biomedical and Clinical Research, Medical University-Pleven, 1 Kliment Ohridski str., 5800 Pleven, Bulgaria; Phone: +359 64 884 304; e-mail: jbcr@mu-pleven.bg

Copyright© 2008 by the Journal of Biomedical and Clinical Research, Medical University-Pleven. All rights reserved.

Journal of Biomedical and Clinical Research

Vol. 14, No. 2, Suppl. 1, 2021

In Memoriam Professor Leon Mitrani:

Scientific Conference in Honour of his 100th Anniversary 16th December 2021

ABSTRACTS

Journal of Biomedical and Clinical Research

Vol. 14, No. 2, Suppl. 1, 2021

In Memoriam Professor Leon Mitrani: Scientific Conference in Honour of his 100th Anniversary

Organized by the Institute of Neurobiology, Bulgarian Academy of Sciences 16th December 2021, Institute of Neurobiology, Sofia

> **Organizing committee** Chair: Assoc. Prof. Dr. Katerina Stambolieva

Members:

Prof. Dr. Reni Kalfin Prof. Assoc. Prof. Dr. Galya Stavreva Assoc. Prof. Dr. Daniela Pechlivanova Assist. Prof. Dr. Milena Mihaylova Assist. Prof. Dr. Polina Mateeva

Scientific committee Chair: Assoc. Prof. Dr. Nadejda Bocheva

Members:

Dr. Juliana Yordanova Prof. Dr. Jana Tchekalarova Prof. Dr. Nikolay Lazarov Assoc. Prof. Dr. Maria Antonova Secretary Svilena Gancheva -In Memoriam Professor Leon Mitrani: Scientific Conference in Honour of his 100th Anniversary, 16th Dec. 2021

Contents

Vol. 14, No. 2, Suppl. 1, 2021

- 7 Coupling Between Slow Waves and Sleep Spindles During Slow Wave Sleep in Humans Depends on Functional Brain States Before Sleep Vasil Kolev, Juliana Yordanova
- 7. Translation Invariance of Visual Object Recognition *Ivan Vankov*
- 8 Motor Theta Oscillations in Aging Juliana Yordanova, Vasil Kolev
- 8 The Stress of Covid-19 and the Development of Fears *Silvia Koumanova*
- 9 Effects of Age and Cue Number in Arbitrary Classification Tasks Bilyana Genova, Simeon Stefanov, Nadejda Bocheva, Miroslava Stefanova
- 9 Comparison of Basic Visual Functions Between Children And Adolescents with Developmental Disorders and Typical Development Milena Slavcheva Mihaylova, Eleonora Encheva-Stoykova, Ivan Hristov, Bilyana Genova, Kalina Racheva, Nadejda Bocheva
- 10 Fixation Stability and Gaze Allocation Area Reflect Differences Among Asd, Adhd, and Dd Groups According to the Task Requirements Nadejda Bocheva, Miroslava Stefanova, Tsvetalin Totev, Simeon Stefanov, Milena Mihaylova
- 11 Stress and Mental Disorders in the Covid-19 Pandemic Yordan Tankovski
- 11 Development of Auditory Attention in Children and Adolescents: Behavioral Study Plamenka Nanova, Vasil Kolev, Juliana Yordanova
- 12 Influence of Schoolbag Load on the Static Postural Stability of 7-Year-Old Children Dragomir Ivanov, Katerina Kirilova, Yoanna Yordanova, Katerina Stambolieva
- 13 Age-Dependent Metabolic Effects in Rats with Pinealectomy Petya Ivanova, Jana Tchekalarova, Lidia Kortenska, Petar Hrischev, Katerina Georgieva
- 13 Neurological Complications After Covid-19 Infection Thomas Kedras, Angeliki Moutafi, Elena Dzhambazov, Daniela Pechlivanova
- 14 In Vitro Evaluation of The Effects of Newly Synthetized Hydrazones and 1,3,4-Oxadiazoles Derivatives on Isolated Rat Brain Synaptosomes and Human Recombinant Mao-B Enzyme Valentin Karabeliov, Reni Yakimova, Violina Stoyanova, Boris Petrov, Magdalena Kondeva-Burdina
- 14 Analgesic Effects of Valorphin Peptide Analogs with Unnatural Cyclic Amino Acids in Rats Borislav Assenov, Petar Todoro, Elena Dzhambazova, Polina Mateeva, Daniela Pechlivanova
- 15 Melatonin Protective Effect Against Aβ- Induced Neurotoxicity and Memory Impairment in Icv Aβ1-42 Rat Model with Pinealectomy Irina Georgieva, Veselina Uzunova, Vanya Milanova, Sonia Apostolova, Jana Tchekalarova, Zlatina Nenchovska, Natasha Ivanova, Tzveta Dimitrova, Lidia Kortenska, Rumiana Tzoneva
- 16 Effect of Prednisolone and Imatinib Mesylate on Amiodarone-Induced Pulmonary Fibrosis in Rats Evgenia Tsolova, Plamen Krastev, Venka Tsankova, Galya Stavreva
- 16 Agomelatine Attenuates Behavioral Changes in Two Rat Models of Sporadic Alzheimer's Disease Kalina Ilieva, Milena Atanasova, Jana Tchekalarova

- 17 Piromelatine Treatment Restores Impaired Sleep-Wake Cycle and Sleep Architecture Through Melatonin Receptors in Male and Female Offspring Rats with Prenatal Stress Natasha Ivanova, Lidia Kortenska, Jana Tchekalarova
- 18 Decreased Expression of Bdnf in the Hippocampus After Prenatal Stress in Rats Is Associated with Changes in Spatial Memory in a Sex-Dependent Manner: Effect of Piromelatine Natasha Ivanova, Zlatina Nenchovska, Rumyana Mitreva, Milena Atanasova, Maria Bakalova, Penka Petelova, Jana Tchekalarova
- 18 The Role of Age and Melatonin Deficiency on Emotional Status and Level of Oxidative Stress in Experimental Rats Jana Tchekalarova, Zlatina Nenochovska, Tzveta Stoyanova, Veselina Uzunova, Rumiana Tzoneva
- 19 Effect of Obestatin on Maximal Force of Contraction of Excised Frog Heart After Chemical Sympathectomy with 3-Iodothyrosine Bilyana Ilieva, Hristo Gagov, Iliyana Sazdov
- 20 Guide for Authors
- 23 Ethical Guidelines
- 26 Submission Form

COUPLING BETWEEN SLOW WAVES AND SLEEP SPINDLES DURING SLOW WAVE SLEEP IN HUMANS DEPENDS ON FUNCTIONAL BRAIN STATES BEFORE SLEEP

Vasil Kolev, Juliana Yordanova

Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Vasil Kolev Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: vasil_kolev@yahoo.com*

Summary

BACKGROUND. Slow wave activity (SWA) and sleep spindles are fundamental electrophysiological signatures of non-rapid eye movement (NREM) sleep in both humans and animals. Previous observations have revealed that these two sleep events are not independent during thalamo-cortical interactions. Furthermore, co-existent sleep spindles and slow waves have been viewed as a mechanism for offline information processing.

OBJECTIVE. The aim was to test the hypothesis that the coupling between SWA and sleep spindles is associated with offline learning during sleep. For this aim, we explored if the temporal synchronization between slow waves and spindle activity during slow wave sleep (SWS) in humans was modulated by preceding brain states, i.e., by functional activations induced by pre-sleep learning.

METHODS. We activated differentially the left and right hemisphere before sleep by using a lateralized variant of serial response time task (SRTT) and verified these inter-hemispheric differences by analyzing alpha and beta electroencephalographic (EEG) activities during learning. A total of 53 subjects were studied with half of them training the right hemisphere, and the other half training the left hemisphere before sleep. The stability and timing of coupling between positive and negative phases of slow waves and sleep spindle activity during SWS were quantified at multiple electrode locations before during sleep.

RESULTS. Spindle activity was temporally synchronized with both positive (up-state) and negative (down-state) slow half waves. Synchronization of

only the fast spindle activity was laterally asymmetric after learning, corresponding to hemisphere-specific activations before sleep. However, the down state was associated with decoupling, whereas the up-state was associated with increased coupling of fast spindle activity over the pre-activated hemisphere.

CONCLUSION. These observations provide original evidence that (1) the temporal grouping of fast spindles by slow waves is a dynamic property of human SWS modulated by functional pre-sleep activation patterns, and (2) fast spindles synchronized by slow waves are functionally distinct.

TRANSLATION INVARIANCE OF VISUAL OBJECT RECOGNITION

Ivan Vankov

Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Ivan Vankov Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: i.vankov@inb.bas.bg*

Summary

An important characteristic of visual perception is its degree of robustness to spatial transformations of the retinal image. For example, during object recognition, the visual system has to establish the identity of an object when it is projected to varying retinal positions - a capacity known as 'translation invariance' or 'translation tolerance'. Previous attempts to assess the translation invariance of biological vision have produced controversial results. In this work, I will present the findings of a series of behavioural studies with human subjects showing that the objects trained at one retinal location can be recognized at high accuracy rates following translations up to 18 visual degrees. Computational modelling with deep convolutional neural networks also demonstrates extreme translation invariance and highlights some of the mechanisms which may underlie this phenomenon.

MOTOR THETA OSCILLATIONS IN AGING

Juliana Yordanova, Vasil Kolev

Institute of Neurobiology, Bulgarian Academy of Sciences **Corresponding Author**: Juliana Yordanova Bl. 23, Acad. G. Bonchev Str.

Bl. 23, Acad. G. Bonchev S Sofia, 1113 Bulgaria e-mail: jyord@yahoo.de

Summary

BACKGROUND. It has been suggested that a distributed oscillatory system in the brain operating in the theta frequency range plays a major role in coordinating motor actions. In this distributed theta system, a central coordinating ("hubbing") function has been attributed to the medial frontal regions including the supplementary motor area (SMA) and the anterior cingulate cortex (ACC). Also, theta activity at motor cortical regions is closely associated with medial frontal theta activity, possibly in relation with movement regulation, monitoring and control. Aging-related alterations have been demonstrated for medial frontal theta activity, but it remains not well understood if theta oscillations at motor cortical regions change with advancing age in humans.

OBJECTIVE. In the present study, electroencephalographic (EEG) theta (3.5-7 Hz) oscillations were analyzed during motor response generation. The major objective was to explore the effects of human aging on the neurodynamics of motor-related EEG theta activity in sensorimotor tasks. EEG theta oscillations were targeted for analysis during motor response generation.

METHODS. Response-related potentials of young and older adults elicited in auditory and visual fourchoice sensorimotor tasks were analyzed in the timefrequency domain. The phase-locking factor and total power were computed at motor cortical regions contra- and ipsilateral to the movement and at the midline to reflect phase synchronization and power of motor theta oscillations.

RESULTS. Major results demonstrated that in both young and older adults, a pronounced responselocked theta activity (3.5-7 Hz) was generated at premotor, motor and sensorimotor cortical regions contra-lateral to the responding hand. Aging was associated with a decreased lateral asymmetry in the phase synchronization of only the left-hand responses. Also, there was a strong aging-related suppression of theta power at the medial fronto-central region that was specifically enhanced and correlated with response speed only in young adults.

CONCLUSIONS. These results confirm the involvement of theta oscillations in movement generation in relation to action coordination in humans. Also, they show that aging is associated with a dysfunction of the oscillatory theta system for motor action regulation due to a strong suppression of a medial frontal integrating mechanism and an impaired balance in the functional control of responses with the right and the left hand.

THE STRESS OF COVID-19 AND THE DEVELOPMENT OF FEARS

Silvia Koumanova

Centre for Stress Management SOMENSO; Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Silvia Koumanova Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: silvia.koumanova@somenso.com*

Summary

A pandemic of COVID-19 has caused a surge in stress experienced by individuals around the world. This fact has given rise to a number of psychological, neuro-behavioral and clinical studies. It has been recognized that much of society's behavioral and emotional responses to the virus can be viewed in the context of the two main fears - fear of disease (nosophobia) and fear of death (thanatophobia). The assumption that poor mental health may increase susceptibility to infection and transmission of the virus during a pandemic has initiated a number of studies worldwide related to the psychological and physiological components of the disease and related underlying fears. Accordingly, theories about the main fears and the reasons for their emergence have been forwarded. It is argued that an increase in the manifestation of post-traumatic stress leads to increased fear of death, which in some cases further results in personality dysfunction. In view of this etiologic frame, it is proposed to emphasize on stress prevention and those post-stress programs that deal with fears. It is assumed that these focused programs would best manage the mental well-being of individuals. Further research is essential to examine the correlation between mental well-being, the course of the pandemic itself and each country's capacity to counteract it.

EFFECTS OF AGE AND CUE NUMBER IN ARBITRARY CLASSIFICATION TASKS

Bilyana Genova, Simeon Stefanov, Nadejda Bocheva, Miroslava Stefanova

Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Bilyana Genova Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: b.genova@abv.bg*

Summary

Humans often classify objects and events in different categories in everyday life. The category learning process requires acquiring the characteristics (cues) of the category members and learning the associations between the stimuli and the responses in repeated presentation trial by trial.

We aimed to investigate the effects of age and the cue number in classification rules on category learning with arbitrary associated stimuli and responses.

The stimuli consisted of randomly distributed elements (blue, red, yellow, and green spheres or cubes) forming two types of dynamic patterns ("wall" and "cloud") that moved either to the left, to the right, forward, or backward. Thus the stimuli differed by four characteristics: color, motion type, motion direction, and shape of the elements. We performed three category learning experiments which varied by the number of the cues defining classification rules used to separate the stimuli into two categories: in Experiment 1, the classification is determined by the color of the elements, in Experiment 2 - by the combination of direction and type of motion, and in Experiment 3 - by the combination of color, direction, and type of motion. In all experiments, the shape of the elements was irrelevant to the category learning task. Two age groups - 16 young (Md = 22 years) and 17 old (Md=67 years) observers participated in the experiments.

The results show that the response time differs nonsignificantly between the two age groups. For the elderly group, neither the classification rule nor the trial number affects the response time. For the young group, the response time increases with the cue number and decreases with the number of stimulus presentations. However, the decrease is slighter for classification by motion direction and type.

The proportion of correct responses depends on the participant group – the elderly group is less accurate. It varies with the number of cues being least for the combined rule. The performance improves with the trial number differently for the two age groups. The elderly group gives less correct responses than the young one when classifying stimuli differing by color or movement. In contrast, the proportion of correct responses is about the same for the combined rule.

The differences in the two age groups' performances are probably due to the changes in working memory with age and the ability to ignore irrelevant information.

COMPARISON OF BASIC VISUAL FUNCTIONS BETWEEN CHILDREN AND ADOLESCENTS WITH DEVELOPMENTAL DISORDERS AND TYPICAL DEVELOPMENT

Milena Slavcheva Mihaylova¹, Eleonora Encheva-Stoykova², Ivan Hristov¹, Bilyana Genova¹, Kalina Racheva¹, Nadejda Bocheva¹

¹Institute of Neurobiology, Bulgarian Academy of Sciences ²Medical Faculty, Department of Physiology and Pathophysiology, Medical University of Sofia

Corresponding Author:

Milena Slavcheva Mihaylova Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: milenski_vis@abv.bg*

Summary

Developmental disorders such as Autism Spectrum Disorder (ASD), Attention Deficit Hyperactivity Disorder (ADHD), and Developmental Dyslexia (DD) are distinguished by a specific pattern of behavioural and learning difficulties with different core defining features. However, they also share some common deficits in sensory information processing. In order to better understand visual perception peculiarities in these disorders, we compared their results on standardized visual tests assessing basic visual functions like visual acuity, contrast sensitivity, colour vision, and stereo vision. One hundred twelve children and adolescents (8-16 years old, 75 boys and 37 girls) separated into four groups with ASD, ADHD, DD, or typical development (TD) participated in the study. All of the tests were performed while the participants wore their lenses or glasses if they already had a refractive error correction.

The results obtained showed similar results between all the groups participating in the study concerning the near visual acuity, stereo acuity, and colour vision anomalies. However, it was observed that a significantly higher number of participants in the groups with ASD and DD have lowered far visual acuity and decreased contrast sensitivity.

The impaired vision in the groups with ASD and DD could predetermine some of their difficulties in learning and education as well as daily activities and communication problems. Moreover, the diminished contrast sensitivity could potentially be associated with the compromised excitation/inhibition balance reported in ASD (and assumed in DD) and would influence the receptive fields' center/periphery relations.

Acknowledgments: This work was supported by the National Science Fund Bulgaria, contract DN15/6/2017

FIXATION STABILITY AND GAZE ALLOCATION AREA REFLECT DIFFERENCES AMONG ASD, ADHD, AND DD GROUPS ACCORDING TO THE TASK REQUIREMENTS

Nadejda Bocheva¹, Miroslava Stefanova¹, Tsvetalin Totev¹, Simeon Stefanov¹, Milena Mihaylova¹

¹Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Miroslava Stefanova Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: mirad_st@abv.bg*

Summary

Developmental disorders, like Autism Spectrum Disorder (ASD), Attention Deficit Hyperactivity Disorder (ADHD), and Developmental Dyslexia (DD), have different diagnostic features, but all of them have an atypical visual perception and eye movements. In the present study, we examined the fixation stability of gaze in children and adolescents from these groups and a group with typical development (TD) in a contour integration task. Sixteen participants from each group (64 in total) took part in the study. They had to detect a straight contour consisting of aligned Gabor patches placed amongst similar Gabor elements with random orientation. We varied the degree of elements' deviation from the contour (visual noise level) in separate blocks. The gaze position of the participants during the fixation point presentation preceding the stimulus was recorded and analyzed. The eye positions in this interval provide a measure of fixation stability and the ability of the participants to allocate gaze to the most informative parts of the subsequent stimulus presentation.

The eye positions were analyzed by spatial point analysis in which the points patterns were described as a realization of an inhomogeneous Poisson process. It allows estimating the differences in the area occupied by the gaze locations and the effect of the group differences and the noise level on it.

The results show that the gaze positions of the participants from the groups with developmental disorders differ in size and elongation from the group with TD suggesting a greater fixation instability. The visual noise affects all groups' gaze point patterns, implying more individual differences in the choice of the most informative parts of the image. The spread of the gaze distributions correlates negatively with the IQ scores, indicating that participants with lower intellectual abilities have more dispersed gaze positions.

Acknowledgments: This work was supported by the National Science Fund Bulgaria, contract DN15/6/2017.

STRESS AND MENTAL DISORDERS IN THE COVID-19 PANDEMIC

Yordan Tankovski

Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Yordan Tankovski Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria e-mail: tankovskibg@abv.bg

Summary

Nowadays, the fact that most people in Bulgaria underestimate the care for their mental health is being increasingly recognized. Unfortunately, for almost two years, humanity has been in a situation of a Covid-19 pandemic, which puts people's lives in an extreme situation, and mental health care must be a top priority, along with pandemic measures and health care.

This presentation will provide a brief overview of what stress is, the types of stress, and types of stress reactions. The typical mental disorders, mental experiences and consequences for people during the Covid-19 pandemic will be outlined in the context of disease-related stress and anti-pandemic restrictions. Special attention will be paid to three most common disorders (according to ICD 10) during the pandemic - Panic Disorder (PA), Generalized Anxiety Disorder (GAD) and Post-Traumatic Stress Disorder (PTSD). psychological Different specific phenomena (cognitive distortions) in post-Covid symptoms will be highlighted. The reported mental problems are extracted basing on observations of the work of Bulgarian psychotherapists and clinical psychologists, who conduct numerous online consultations.

Another goal of the presentation is to emphasize on the psychological consequences of not treating the described anxiety disorders. Basic measures will be outlined that can be taken to reduce health damage by providing psychological assistance to victims of these disorders. It will be emphasized that anxiety disorders can be managed, but this depends on the acquisition of basic self-help and assistance skills. Finally, several typical cognitive distortions will be described that have been identified in people who appear in force majeure circumstances such as restrictions, negative news, bad messages from politicians, isolation, fear of getting sick with Covid 19, and uncertainty about the future.

DEVELOPMENT OF AUDITORY ATTENTION IN CHILDREN AND ADOLESCENTS: BEHAVIORAL STUDY

Plamenka Nanova, Vasil Kolev, Juliana Yordanova

Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Plamenka Nanova Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: pnanova@yahoo.com*

Summary

BACKGROUND. "Attention is the taking possession of the mind, of one out of several simultaneously possible objects or trains of thought ... It implies withdrawal from some things in order to deal effectively with others" (James W., 1890). "The extent that selective attention skills are relevant for academic foundations and amenable to training. they represent an important focus for the field of education" (Stevens and Bavelier, 2012). OBJECTIVE. The aim of the present work was to study the behavioral performance of children and adolescents during auditory selective attention task. METHOD. In the auditory selective attention task, sequences of two stimuli with different frequency (40% targets and 60% non-targets) were presented randomly to the right and left ear. During the task participants maintained lateralized internal attention either to right- or left-side stimuli in order to produce

either to right- or left-side stimuli in order to produce selective motor responses with the right hand to targets appearing in the attended side. Thus, there were two conditions - attend-right and attend-left, each comprising 4 stimulus types - target-attended, non-target-attended, target-non-attended, non-targetnon- attended. Correct response rate, reaction times (RTs) to targets, and response variance (RV) measured by RT standard deviation and coefficient of variance, were registered and analyzed in 110 normal, healthy, right-handed children and adolescents, divided into four age groups and two gender groups (9-16-yearsold, age groups: 9-10, 11-12, 13-14 and 15-16-yearsold). Effects of age, gender and side of attention were explored.

RESULTS. The rate of correct responses increased

with age and was also significantly higher in the attend-right than attend-left condition. RTs decreased significantly with development and were significantly shorter in the attend-right than attend-left condition. RV decreased significantly with development. There were no significant differences in RV between attend-right and attend-left conditions. In addition, no performance or RT/RV gender differences were found between girls and boys.

CONCLUSIONS. These performance results demonstrate that the mechanisms of auditory selective attention progressively improve with development from 9 to 16 years. Notably, this developmental improvement is equally expressed in males and females. The observation that focusing internal attention to the right side produces faster and more accurate responses as compared to the attendleft responses may reflect effects of lateral asymmetry in sensory and motor activations. Given that righthand responses were required in the present task, the attend-right advantage implies a more efficient processing with unilateral activation of sensory and motor areas in only the left hemisphere, in contrast to the bi-lateral activation in the two hemispheres in the attend-left condition (right sensory and left motor areas). The proposed effects of (non)-lateralized activations persist as early as 9 years of age, do not change with development, and are independent of gender.

INFLUENCE OF SCHOOLBAG LOAD ON THE STATIC POSTURAL STABILITY OF 7-YEAR-OLD CHILDREN

Dragomir Ivanov, Katerina Kirilova, Yoanna Yordanova, Katerina Stambolieva

Institute of neurobiology, Bulgarian Academy of Science

Corresponding Author:

Dragomir Ivanov Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: dkivanov@gmail.com*

Summary

The schoolbags are the preferred choice due to the fact that they distribute the weight evenly on the

spine without overloading only one shoulder. The schoolbag overload has negative effects on the correct posture and leads to neck, shoulder and back pain. In children this overload can lead to both temporary and chronic musculoskeletal problems. It was found that the weight of the load should not exceed 15% of the total body weight of the child carrying it.

The aim of this study is to evaluate the effects of schoolbag load on the maintenance of static postural stability in 7-year-old children from elementary school.

The postural sway was measured by static posturographic system during quiet stance on firm support in six experimental conditions: stance with open eyes; stance with closed eyes without load and the same condition with backpack load with weights 2-3% and 16-17% of the child's body weight. Thirty-six participants took part in the investigation.

The results indicate that postural stability with a backpack weighing about 16-17% of the body weight are significantly different from the conditions without backpack and with 2-3% load. An increase in all postural parameters during standing under load was found, but the increase in those parameters in condition under load about 16-17% was significant compared to two other conditions (no-load and 2-3% load). In this condition, a significant increase in mean sway velocities in both directions (AP and ML) was observed during standing with open and closed eyes, while the mean sway amplitudes increased significantly in standing with eyes closed only. The results suggest that standing with a loaded backpack mostly affects the velocities of postural sway than its amplitudes. This investigation show that the load on the back over 15% of the total body weight of the child increases the postural sway and leads to instability of static upright stance.

Key words: children, backpack weight, postural stability

Acknowledgments: This work was partially supported by the Bulgarian Ministry of Education under the National Research Programme "Young scientists and post-doctoral students".

AGE-DEPENDENT METABOLIC EFFECTS IN RATS WITH PINEALECTOMY

Petya Ivanova¹, Jana Tchekalarova¹, Lidia Kortenska¹, Petar Hrischev², Katerina Georgieva²

¹Institute of Neurobiology, Bulgarian Academy of Sciences ²Department of Physiology, Medical University-Plovdiv

Corresponding Author:

Petya Ivanova Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: ivanova.petya91@gmail.com*

Summary

The pineal gland is suggested to be an important area involved in programming of fertility, growth, aging and death of mammal via the released hormone melatonin. The aim of the present study was to ascertain the effect of melatonin deficit on several metabolic parameters, closely associated with aging process, at certain stages of ontogenesis. Sham and rats with pinealectomy, surgered at age of 3-, 14- and 18-months, respectively, were tested two months later. Age-related decline was demonstrated in sham-operated rats as decreased skeletal muscle strength, food intake as well as worsened calorimetric parameters. Pinealectomy decreased the maximal time to exhaustion, food intake, oxygen consumption, CO2 production and energy expenditure during the dark period in the youngest rat group. In addition, melatonin deficit elevated blood pressure (systolic, diastolic and mean arterial pressure) as well as impaired serum glucose profile in 3-month-old rats with pinealectomy. In conclusion, the present study suggested that melatonin deficit accelerate aging associated with crucial physiological, metabolic and biochemical parameters in young adult rats.

Key words: Pinealectomy, Aging, metabolism, Rats

Acknowledgments: This work was supported by the National Science Fund of Bulgaria (research grant # DN 12/6 and # ΚΠ-06-ΠΗ-41/1, 2020).

NEUROLOGICAL COMPLICATIONS AFTER COVID-19 INFECTION

Thomas Kedras¹, Angeliki Moutafi¹, Elena Dzhambazova¹, Daniela Pechlivanova^{1,2}

¹Medical Faculty, Sofia University, St Kliment Ohridski ²Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Angeliki Moutafi 1, Koziak Str. Sofia, 1407 Bulgaria *e-mail: angeliki.moutafi@outlook.com*

Summary

INTRODUCTION

COVID-19 pandemic has affected global health at an unprecedented rate, infecting 255 million people, while leading to death more than 5 million. The coronavirus subtype SARS-CoV-2, is an mRNA virus, which even though attacks primarily the cardiovascular and respiratory systems, it is a well-known fact that neurological manifestations are not being missed. AIM

The aim of this literature review is to summarize the latest research data available and provide healthcare workers with an up-to date overview, regarding the neurological complications. MAIN BODY

SARS-CoV-2 has a great affinity to the ACE2 receptor and it is supported that its migration, and subsequently, infection is facilitated by the transmembrane protease serine 2 (TMPRSS2) which helps with the viral binding to those receptors. ACE2 receptors are expressed primarily on the pulmonary alveolar cells, while extrapulmonary in the vascular endothelial cells of the GI tract, heart and brain and on the neurons and glial cells. Therefore, apart from respiratory (dry cough, dyspnea) and cardiovascular symptoms (blood clots, ischemia), neurological features are also manifested e.g. anosmia, ageusia, myalgias. More specifically, encephalitis, meningoencephalitis and seizures, are common complications which follow COVID-19 infection. Intracerebral hemorrhage and hemophagocytic lymphohistiocytosis are conditions, that even though have different pathophysiological mechanisms, both are life threatening. Regarding demyelinating disorders, Guillain Barré syndrome

may be presented even after primary infection while transverse myelitis and acute disseminated encephalomyelitis are not often presented clinical entities. Multiple Sclerosis (MS), Alzheimer's disease (AD), and Parkinson's disease (PD), which are classified as neurodegenerative diseases, are suggested to be linked both as risk factors and long term complications due to neuronal damage and loss. CONCLUSION

In view of the above, COVID-19 can affect the human organism unpredictably, having complications ranging from mild to severe. However, there is not enough information on whether or not SARS-Cov-2 infection could accelerate AD, PD, and MS presentation in the long term and more research needs to be done.

IN VITRO EVALUATION OF THE EFFECTS OF NEWLY SYNTHETIZED HYDRAZONES AND 1,3,4-OXADIAZOLES DERIVATIVES ON ISOLATED RAT BRAIN SYNAPTOSOMES AND HUMAN RECOMBINANT MAO-B ENZYME

Valentin Karabeliov^{1,2}, Reni Yakimova¹, Violina Stoyanova², Boris Petrov², Magdalena Kondeva-Burdina¹

¹MU-Sofia, Faculty of Pharmacy, Department "Pharmacology, Pharmacotherapy and Toxicology", Laboratory "Drug metabolism and drug toxicity" ²MU-Sofia, Faculty of Pharmacy, Department "Chemistry"

Corresponding Author:

Boris Petrov 15, Acad. Ivan Geshov blvd. Sofia, 1431 Bulgaria *e-mail: bobi.stoyanov@abv.bg*

Summary

This study evaluates the *in vitro* effects of two series compounds: 1,3,4-oxadiazoles (**3a-k**) and hydrazones derivatives (**5a-n**) on isolated rat brain synaptosomes and human recombinant MAO-B enzyme (hMAO-B). In order to evaluate drug likeness of these newly synthesized molecules, in-silico ADME evaluation was carried out.

On isolated rat brain synaptosomes, administered alone at concentration 40 μ M, both series revealed weak statistically significant neurotoxic effects, compared to the control (non-treated synaptosomes). In conditions of 6-hydroxydopamine-induced oxidative stress, all the compounds showed statistically significant neuroprotective effects, by preserving the synaptosomal viability and the level of reduced glutathione, compared to the toxic agent. On the activity of human recombinant MAO-B enzyme (hMAO-B), all derivatives didn't show statistically significant inhibitory effect, compared to

the control (pure hMAO-B). Of the series of substances studied, only **3a** and **3d** showed a statistically significant inhibitory effect on the activity of human recombinant MAO enzyme (hMAOB). The inhibitory activity of the substances is close to that of Selegiline.

2-(4-Chlorophenyl)-5-phenyl-1,3,4-oxadiazole **3a** inhibits enzyme activity by 44%, 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-1H-indole **3d** inhibits by 46%, and Selegiline (classic MAOB inhibitor) reduces enzyme activity by 55% compared to the control (pure hMAOB). It is worth mentioning that 2,5-substituted 1,3,4-oxadiazoles would represent a productive matrix for the development of new class of MAOB inhibitors.

ANALGESIC EFFECTS OF VALORPHIN PEPTIDE ANALOGS WITH UNNATURAL CYCLIC AMINO ACIDS IN RATS

Borislav Assenov^{1,2}, Petar Todorov³, Elena Dzhambazova², Polina Mateeva¹, Daniela Pechlivanova^{1,2}

¹Institute of Neurobiology, Bulgarian Academy of Sciences ²Faculty of Medicine, Sofia University "St. Kl. Ohridski" ³University of Chemical Technology and Metallurgy, Sofia

Corresponding Author:

Borislav Assenov Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: b.assenov@abv.bg*

Summary

Introduction. VV-hemorphin-5 (Valorphin or V1) is an endogenous peptide that shows a significant antinociceptive effect. The subject of the present study is two newly synthesized valorphin peptides (C5-V and C6-V) with unnatural cyclic amino acids. Our previous experimental data indicate that intracerebroventricular injection of VV-hemorphin-5 and its analogs causes dose-dependent antinociceptive effects in a formalin test. The aim of the present study was to evaluate the effects of intraplantar injected V1, C5-V, and C6-V on the Carrageenan-induced hyperalgesia as compared to the reference anti-inflammatory and analgesic drug Indomethacin.

Methods. The Fmoc chemistry (solid-phase peptide synthesis) was used to synthesize two new analogues of Valorphin – C5-V (Val-Val-Tyr-**AC5C**-Trp-Thr-Gln-NH₂) and C6-V (Val-Val-Tyr-**AC6C**-Trp-Thr-Gln-NH₂). Based on previous data, all peptides were injected intraplantar in the right hind paw at a dose of 50 μ g / 5 μ l per rat, 5 minutes before the irritant. The inflammatory process was caused by injecting 100 μ l of carrageenan 1% into the same hind paw. The nociceptive effect was assessed using a paw pressure analgesimeter at 1, 3, and 4 hours after carrageenan injection.

Results. Experimental data show that carrageenaninduced inflammation lowers the mechanical pain threshold in the injected paw. V1 showed a significant analgesic effect in the ipsilateral paw at 1st, 3rd, and 4th hours after carrageenan injection. Its effect was higher compared to the reference Indomethacin. C5-V showed a significant antinociceptive effect, comparable to that of V1 after the 1st hour. C6-V also showed a considerable analgesic effect, and its effect was comparable to those of Indomethacin but lower to the V1 induced antinociception.

Conclusion. The results demonstrated a significant and long-lasting local antinociceptive effect of Valorphin in the paw pressure test, while its synthetic analogs C5-V and C6-V showed a lower but comparable to the reference antinociceptive effect of Indomethacin.

Key words: valorphin peptides, nociception, rats **Acknowledgments:** *This work was supported by the Sofia University "St.Kliment Ohridski", contract No.80-10-4 /2021.*

MELATONIN PROTECTIVE EFFECT AGAINST Aβ- INDUCED NEUROTOXICITY AND MEMORY IMPAIRMENT IN ICV Aβ1-42 RAT MODEL WITH PINEALECTOMY

Irina Georgieva¹, Veselina Uzunova¹, Vanya Milanova¹, Sonia Apostolova¹, Jana Tchekalarova², Zlatina Nenchovska², Natasha Ivanova², Tzveta Dimitrova², Lidia Kortenska², Rumiana Tzoneva¹

¹Institute of Biophysics and Biomedical Engineering, Bulgarian Academy of Sciences ²Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Rumiana Tzoneva Bl. 21, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: tzoneva@bio21.bas.bg*

Summary

Melatonin is a multifunctional molecule and its role in the regulation and protection of the central nervous system is well documented. Therefore, melatonin was proposed as a possible therapeutic agent for reducing the severity of Alzheimer's disease (AD), a progressive neurodegenerative disease characterized by cognitive decline and memory dysfunction. In addition, melatonin has a higher preference in comparison to other available antioxidants in the therapeutic strategy of AD due to its inhibitory effect on amyloid beta $(A\beta_{1-42})$ accumulations, low toxicity effect in high doses, and readily passage through blood brain barrier. Pinealectomy appears to be a good model to study melatonin deficiency and pathway dysfunction in the pathogenesis of neurodegenerative diseases.

Hence, we investigated the effect of melatonin on a reduction of the neurotoxicity expressed by Ab and γ -secretase production and memory loss in icv A $\beta_{1.42}$ rat model of AD combined with melatonin deficiency. Animals (ten weeks old male Sprague-Dawley rats) undergone icv infusion of A $\beta_{1.42}$ /vehicle, were

injected with melatonin (50 mg/kg, intraperitoneally (i.p.)/vehicle about two hours before the onset of the dark phase for 40 days. Object recognition test (ORT) and radial arm maze test (RAM) were used to explore cognitive functions. Amyloid beta and gamma secretase levels were measured in the frontal cortex (FC) and the hippocampus. Pinealectomy did not exacerbate associative and spatial memory of $A\beta_{1,42}$ -treated rats in ORT and RAM, respectively. Chronic melatonin administration corrected impaired cognitive performance in both AD model and AD model combined with melatonin deficit. The $A\beta_{1,42}$ secretion was significantly increased compared to sham group in the FC and the hippocampus in the $A\beta_{1,42}$ rats with pinealectomy which effect was compensated by melatonin treatment. The main enzyme which is responsible for A β accumulation - γ -secretase, was elevated in the Aβ-pin groups and the treatment with melatonin lowers the levels to the control values. The present data suggest pinealectomy+A $\beta_{1,42}$ rat model as a more reliable model of AD. Melatonin could have a substantial potential to suppress $A\beta_{1,42}$ accumulation in the brain and concomitant memory impairment **Key words**: icvA $\beta_{1,42}$; melatonin; A β , γ -secretase, memory loss; rat

Acknowledgments: This work was supported by projects FNI № KP-06-H31/16.

EFFECT OF PREDNISOLONE AND IMATINIB MESYLATE ON AMIODARONE-INDUCED PULMONARY FIBROSIS IN RATS

Evgenia Tsolova, Plamen Krastev, Venka Tsankova, Galya Stavreva

Dept. Pharmacology and Toxicology, Faculty of Pharmacy, Medical University – Pleven

Corresponding Author:

Galya Stavreva 1, Saint Kliment Ohridski Str. Pleven, 5800 Bulgaria *e-mail: drstavreva@yahoo.com*

Summary

The experimental model of amiodarone (AM)induced lung toxicity is one of the relevant models to study pulmonary fibrosis. AM administered intratracheally induces inflammatory response, activation of fibroblasts, and subsequent fibrosis. We tested the anti-inflammatory and antifibrotic effect of imatinib mesylate (IM) and prednisolone (PR) against AM-induced pulmonary fibrosis in a rat model.

Methods: The study was carried out on 72 male Wistar rats divided into six treatment groups:

control; AM group – treated intratracheally with AM on days 0 and 2; PR1 group – treated with AM and PR 10 mg/kg orally from day 0 to day 10; PR2 group – treated with AM and PR from day 10 to day 28; IM1 group – treated with AM and IM 50 mg/kg orally from day 0 to day 10; IM2 group – treated with AM and IM from day 10 to day 28. Pulmonary fibrosis was assessed by measuring hydroxiproline (HP) and collagen (COL) content in lung homogenate (LH) and histopathologically on day 28 after AM administration.

Results: AM resulted in significantly increased HP and COL content in LH on day 28. The content of HP and COL in LH of animals, treated with AM and PR, given after day 1 and AM plus IM, given after day 10 were decreased compared to AM alone on day 28 (p<0.05).

Intratracheal AM led to moderate interstitial and perivasal fibrosis, thickening of interstitial spaces and cellular infiltration; these damages were attenuated by above-mentioned dosing regimens.

Conclusion: The results obtained from our study showed that pulmonary fibrosis was attenuated by IM, given during fibrotic phase and by PR, given during inflammatory phase.

Key words: amiodarone, imatinib mesylate, lung fibrosis, prednisolone

AGOMELATINE ATTENUATES BEHAVIORAL CHANGES IN TWO RAT MODELS OF SPORADIC ALZHEIMER'S DISEASE

Kalina Ilieva¹, Milena Atanasova¹, Jana Tchekalarova²

¹Department of Biology, Medical University of Pleven ²Institute of Neurobiology, Bulgarian Academy of Sciences

Corresponding Author:

Kalina Ilieva 1, Saint Kliment Ohridski Str. Pleven, 5800 Bulgaria *e-mail: kalina.st.ilieva@gmail.com*

Summary

In two consequential studies we have investigated the effect of agomelatine against behavioral, biochemical and histological impairments in streptozotocin (STZ) and amyloid beta (A β) induced model of ADs in male rats. The agomelatine, which is a melatonin MT1 and MT2 agonist and 5-HT2C antagonist, was injected intraperitoneally for 30 days, at a dose of 40mg/kg in both studies. We measured the anxiety levels, spatial memory errors and anhedonia by using the following methods: Elevated Plus Maze (EPM), Radial Arm Maze (RAM), Saccharine Preference Test (SPT). The treatment with Aß significantly decreases the number of entries in to the aversive open arms of the EPM and the time spent inside of them for the $A\beta$ group. Reduced anxiety-like behaviour, tested with the EPM, was observed and in the study with STZ-treated rats as they spent longer time on the open arms and made more entries, compared to the group with sporadic AD. Agomelatine corrected anxiety behavior of Aß rats as the treated animals enter the open arms more often and spend significantly longer time in there. Administration of agomelatine decreased working memory errors during the last two sessions in RAM in the model with izvSTZ-induced AD. Testing the Aβ-induced animals showed alleviation in the spatial memory impairment only during the last session. The SPT helped us to investigate the level of anhedonia as a marker of depressive-like behaviour, which is considered as a comorbid condition in AD. Preference for sweet solutions has decreased after icv infusion of A β and STZ. The A β -veh and STZ-veh groups demonstrated anhedonia, compared to the sham-veh group. Agomelatine reversed depressive-like response induced by icv infusion of both toxins, compared to the control groups and increased the preference to saccharine. The ability of Ago to alleviate behavioral symptoms observed in the two models of sporadic ADs suggests that this melatonin alternative can be considered a promising adjuvant in this disease.

PIROMELATINE TREATMENT RESTORES IMPAIRED SLEEP-WAKE CYCLE AND SLEEP ARCHITECTURE THROUGH MELATONIN RECEPTORS IN MALE AND FEMALE OFFSPRING RATS WITH PRENATAL STRESS

Natasha Ivanova^{1,2}, Lidia Kortenska¹, Jana Tchekalarova¹

¹Institute of Neurobiology, Bulgarian Academy of Sciences ²Medical University Sofia, Faculty of Medicine

Corresponding Author:

Natasha Ivanova Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: n.ivanova@inb.bas.bg*

Summary

The prenatal stress (PNS) model in rodents is known to impair circadian rhythms of sleep/wake cycles. Piromelatine (Pir) is a novel melatonin analog designed for the treatment of insomnia. The present study was aimed to explore the effects of Pir and the role of melatonin receptors on PNS-induced impaired sleep architecture and rhythmicity of sleep/ wake cycle and in a rat model of PNS in male and female offspring. The electroencephalographic (EEG) and electromyographic (EMG) recording for up to 24 h were analyzed under basal conditions (12:12 LD cycle+vehicle), PNS+vehicle, PNS+Pir (20 mg/kg/day for 7 days) and PNS+Pir+Luzinodol (Luz) (10 mg/kg) conditions. The rats exposed to PNS were characterized by disturbed sleep/wake cycle associated with diminished NREM sleep, elevated REM sleep and wake pattern in a sexdependent manner. Piromelatine corrected impaired sleep architecture and sleep-wake cycle, while the melatonin (MT) receptor antagonist Luz suppressed the beneficial effect of this melatonin analog. Our findings suggest that Pir can exert sex-dependent beneficial effects on PNS-induced impaired sleep architecture and rhythmicity of sleep/wake cycle via MT receptors.

Key words: prenatal stress, Piromelatine, sleep/ wake cycle, sleep architecture, MT receptors

Acknowledgments: This work was supported by the National Science Fund of Bulgaria (research grant # DN 12/6, KΠ-06-H21/10, KΠ-06-ΠH-41/1, 2020).

DECREASED EXPRESSION OF BDNF IN THE HIPPOCAMPUS AFTER PRENATAL STRESS IN RATS IS ASSOCIATED WITH CHANGES IN SPATIAL MEMORY IN A SEX-DEPENDENT MANNER: EFFECT OF PIROMELATINE

Natasha Ivanova^{1,3}, Zlatina Nenchovska¹, Rumyana Mitreva¹, Milena Atanasova², Maria Bakalova³, Penka Petelova³, Jana Tchekalarova¹

¹Institute of Neurobiology, Bulgarian Academy of Sciences ²Department of Biology, Medical University of Pleven ³Medical University Sofia, Faculty of Medicine

Corresponding Author:

Natasha Ivanova Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: n.ivanova@inb.bas.bg*

Summary

Prenatal stress (PNS) leads to cognitive alterations mainly observed later in life. We investigated the effects of piromelatine, a mixed (MT1) and MT2 and serotonin (5-HT) 1A/1D receptors agonist and 5-HT2B receptor antagonist, on hippocampusdependent spatial memory and concomitant changes in BDNF levels in male and female rat offspring with a history of prenatal stress. Different stressors were applied on pregnant female rats, which started on the 7th gestational day and continued until birth. At the age of 60-days, offsprings were injected every day with piromelatine/vehicle at a dose of 20 mg/ kg for 21 days. Hippocampus-dependent spatial memory was explored in the radial arm maze test (RAM). The BDNF level in the hippocampus was measured by ELISA. A deficit in spatial memory was detected in male but not female rats with prenatal stress compared to their matched controls. Chronic piromelatine treatment corrected the altered memory function and decreased both working and double memory errors in the last trial of the RAM task in the male PNS offspring. The prenatal stress decreased

the expression of BDNF in the hippocampus of male offspring with prenatal stress while female offspring were not affected. The melatonin analogue elevated BDNF in male offspring. The present findings suggest that the novel drug piromelatine could positively affect spatial memory via BDNF in the hippocampus in a sex-dependent manner in rats with a history of prenatal stress.

Key words: prenatal stress, Piromelatine, memory deficit, BDNF

Acknowledgments: This research was supported by the Bulgarian National Science Fund - grant No. KII 06-H21/10.

THE ROLE OF AGE AND MELATONIN DEFICIENCY ON EMOTIONAL STATUS AND LEVEL OF OXIDATIVE STRESS IN EXPERIMENTAL RATS

Jana Tchekalarova¹, Zlatina Nenochovska¹, Tzveta Stoyanova¹, Veselina Uzunova², Rumiana Tzoneva²

¹Institute of Neurobiology, Bulgarian Academy of Sciences

²Institute of Biophysics and Biomedical Engineering, Bulgarian Academy of Sciences

Corresponding Author:

Jana Tchekalarova Bl. 23, Acad. G. Bonchev Str. Sofia, 1113 Bulgaria *e-mail: janetchekalarova@gmail.com; jt.chekalarova@inb.bas.bg*

Summary

The pineal gland has a crucial role on development and aging process via the released hormone melatonin. In the present study, we aimed to evaluate the influence of melatonin deficit at different aging stages on behavior and oxidative stress level in the hippocampus of rats operated 3-, 14- and 18-month-old. The anxiety and depressive-like behavior was elevated in age-depended manner in the sham-operated rats. The removal of the pineal gland induced increased locomotor activity in the open field of the youngest group and provoked depressive-like behavior in both rats operated 3- and 14-month-old, respectively. The superoxide dismutase (SOD) activity was enhanced in the sham-operated 14-month-old rats compared to the matched younger rats while there were no changes in the level of glutathione (GSH) among different ages in the sham groups. The lipid peroxidation was elevated in mature and aged rats compared to the youngest sham rats. Pinealectomy suppressed the activity of SOD and GSH and increased malondialdehyde level in the hippocampus compared the matched sham 14-month-old rats. In conclusion, our results suggest that behavioral responses associated with emotional status and oxidative stress level are vulnerable to changes in young and mature rats with pinealectomy while melatonin deficiency has no impact in elder animals.

Key words: Age, Melatonin, behavior, oxidative stress, rat.

Acknowledgments: This work was supported by the National Science Fund of Bulgaria (research grant # DN 12/6 and # **ΚΠ-06-ΠΗ-41/1, 2020**).

EFFECT OF OBESTATIN ON MAXIMAL FORCE OF CONTRACTION OF EXCISED FROG HEART AFTER CHEMICAL SYMPATHECTOMY WITH 3-IODOTHYROSINE

Bilyana Ilieva^{1*}, Hristo Gagov¹, Iliyana Sazdova¹

¹Department of Human and Animal Physiology, Faculty of Biology, Sofia University "St. Kliment Ohridski"

Corresponding Author:

Bilyana Ilieva 8, Dragan Tsankov Blvd. Sofia, 1164 Bulgaria *e-mail: b.ilieva@biofac.uni-sofia.bg*

Summary

The aim of this study is to investigate the positive inotropic effect of obestatin in the presence of 3-iodothyrosine (3-IT) in heart preparations of Pelophylax ridibundus frog. It is known that, the myocardial β -adrenoreceptors and protein kinase A downstream targets are responsible for the observed positive inotropic effect of obestatin. The application of obestatin (1 nmol/l and 100 nmol/l) significantly enhances the force of contraction of excised and cannulated frog hearts. This effect was completely blocked in the presence of 3-IT, a selective inhibitor of tyrosine hydroxylase that is the rate-limiting enzyme in the catecholamine biosynthesis chain. 3-IT was injected into the dorsal lymph sac 1 hour before isolation and cannulation of the heart at a dose of 0.02 mg/g b.w. It is concluded that, the treatment with 3-IT inhibits adrenaline synthesis and in this way abolishes adrenaline mediated positive inotropic effect of obestatin.

Key words: Obestatin, Cardiac, Hormone, Autonomic nervous system, In vitro, Sympathectomy **Acknowledgments:** This work was supported by grant № 80-10-108/25.03.2021 from Sofia University "St. Kliment Ohridski".

GUIDE FOR AUTHORS

Journal of Biomedical and Clinical Research (JBCR) is a peer reviewed, multidisciplinary scientific journal of Medical University - Pleven, open to scientists from all fields of medicine and related areas.

JBCR publishes reviews, original articles and communications in the field of biomedical, clinical, public health and health care research, medical education, case reports, written in English. **JBCR** is published two times per year.

JBCR follows uniform requirements for manuscripts submitted to biomedical journals, developed by the International Committee of Medical Journal Editors (www.ICMJE.org).

All human and animal studies reported should have been approved by the Institutional Ethical Committee in accordance with the ethical criteria concerning experiments involving human beings or animals.

Submission of an article implies that the work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis), that it is not under consideration for publication elsewhere, that its publication is approved by all authors, and tacitly or explicitly - by the responsible authorities where the work was carried out. Authors should take responsibility for the integrity of the work as a whole, from inception to published article.

Authors are requested to disclose any possible conflicts of interest. Affiliation of authors submitting manuscripts in their role as members of industrial or government organizations may constitute sufficient disclosure. Academic, nonprofit, consulting, advocacy, and all other authors are asked to disclose in a footnote all sources of financial and material support for the studies and manuscripts submitted.

Each manuscript must be accompanied by a submission form, sighed by each author. This should include the name and mailing address, telephone and fax numbers of the corresponding author, who is responsible for communications with the other authors about revisions and final approval of the proof.

1. Submission of papers

Two copies of the manuscripts (on one side of the paper only), written in English, accompanied by a CD (labeled with the file name) should be submitted to: Radoslav Hristov

Editorial Office of *Journal of Biomedical and Clinical Research* Medical University 1, St. Kliment Ohridski, Str. 5800 Pleven, Bulgaria Tel. 064/884 110 *e-mail: jbcr@mu-pleven.bg*

2. Presentation of manuscript

Manuscripts should be submitted as a text file saved in Microsoft Word 2003. The entire text should be typed, using font Times New Roman, 12 points; double spaced and wide (2.5 cm) margins on standard A4; left aligned. Please avoid full justification, i.e., do not use a constant right-hand margin). Headings should be bold; only the first letter of the first word should be capitalized, left aligned. Divide your article into clearly defined sections. Do not use bullets for subsections. They should be numbered 1.1 (then 1.1.1, 1.1.2, ...), 1.2, etc. Any subsection may be given a brief heading. Each heading should appear on its own separate line. Ensure that each new paragraph is clearly indicated (use "Enter" only at the end of paragraphs). There should be no bank lines between paragraphs. Do not use tabs or spaces to create indents. Use the Symbol font for symbols and special characters. Use decimal points, not commas (e.g. 5.06); use a space for thousands (10 000 and above). Present tables and figure captions on separate pages at the end of the manuscript. Number all pages consecutively with Arabic numerals on the right timse reports - 6.

Each manuscript should follow the following sequence: title page; summary; text (Introduction, Materials/Patients and Methods, Results, Discussion, Conclusions); references; tables (table complete with title and footnotes on a separate page); figure legends.

Nonconforming manuscripts (e.g., improper format, excessive length) may be returned for technical revision prior to undergoing peer review.

Title (first) page

The title should be concise and informative.

Names of author(s). Full first name and last name, and middle-name initials of all authors.

Affiliations. State the author's (authors') affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript digit (number) immediately after the author's name and in front of the appropriate address. The name and institution of the first author (and other authors of the same institution) should not bear any number.

Corresponding Author. Phone, address, e-mail.

Acknowledgements. Authors should identify financial support for research, including information on grants received. Technical help, critical reviews of the manuscript and financial or other sponsorship should be acknowledged.

Second page

Summary. A concise and factual summary is required (not more than 200 words). The abstract should briefly state the purpose of research, principal results and major conclusions. References should therefore be avoided, but if essential, they are to be cited in full. Abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

Key words. Immediately after the abstract, provide a maximum of 5 keywords, strictly referring to the Medical Subject Headings list of the Index Medicus (http://www.nlm.nih.gov/mesh/MBrowser.html).

Third page

Title, names of authors and summary in Bulgarian should be given for online version of the journal.

Next pages

Text should carry the following obligatory sections: Introduction.

Materials/Patients and Methods. Provide sufficient detail to allow the work to be reproduced. Methods already published should be indicated by a reference: only relevant modifications should be described.

Results. Results should be described clearly and concisely. All measurements should be reported in International System of Units (SI). Statistical significances should be shown along with the data in the text, as well as in tables and figures. Provide exact p-values, with three decimal places.

Discussion. This should explore validity and significance of the results from the work, and results considered in the context of results in other trials reported in the literature.

Conclusions.

References. JBCR follows the International Committee of Medical Journal Editors Uniform Requirements for Manuscripts Submitted to Biomedical Journals

(http://www.nlm.nih.gov/bsd/uniform_requirements.html). References should be numbered sequentially [in square brackets] in the text and listed in the same numerical order in the reference section. Do not put a period after the reference number. Separate reference number and (last) name of the first author by one space only. All references should be complete and accurate. Unpublished results and personal communications are not recommended in the reference list but may be mentioned in the text. Citation of a reference as "in press" implies that the item has been accepted for publication. Journal titles should conform to the present Index Medicus abbreviations

(http://www.ncbi.nlm.nih.gov/sites/entrez?db=journals)

Examples from http://www.nlm.nih.gov/bsd/uniform_requirements.html

Standard journal article Provide names of all authors when there are six or fewer, if more - list the first six authors followed by et al.

Halpern SD, Ubel PA, Caplan AL. Solid-organ transplantation in HIV-infected patients. N Engl J Med. 2002;347(4):284-7.

Issue with supplement

Glauser TA. Integrating clinical trial data into clinical practice. Neurology. 2002;58(12 Suppl 7):S6-12. *Issue with no volume*

Banit DM, Kaufer H, Hartford JM. Intraoperative frozen section analysis in revision total joint arthroplasty. Clin Orthop. 2002;(401):230-8.

Author(s) and editor(s)

Breedlove GK, Schorfheide AM. Adolescent pregnancy. 2nd ed. Wieczorek RR, editor. White Plains (NY): March of Dimes Education Services; 2001.

Chapter in a book

Meltzer PS, Kallioniemi A, Trent JM. Chromosome alterations in human solid tumors. In: Vogelstein B, Kinzler KW, editors. The genetic basis of human cancer. New York: McGraw-Hill; 2002. p. 93-113. *Journal article on the Internet*

Abood S. Quality improvement initiative in nursing homes: the ANA acts in an advisory role. Am J Nurs [serial on the Internet]. 2002 Jun [cited 2002 Aug 12];102(6):[about 3 p.]. Available from: http://www.nursingworld.org/AJN/2002/june/Wawatch.htm

Tables and Figures should be given on separate sheets with indications on the left hand margin to the text, as to the appropriate placement (highlighted in yellow - Table 1.). Tables and figures should be numbered consecutively in order of their citation in the body of the text, with Arabic numerals. Supply the word "Table" or "Figure", followed by its respective number and a brief title. The footnotes should contain all the necessary explanations of symbols and findings, and statistics. For the footnotes use the following symbols: *, †, ‡, §, ††, ‡‡,...

Data Tables should be submitted in Microsoft Word table format. Tables should not contain vertical lines. Give each column a short heading. Be sparing in the use of tables and ensure that the data presented in tables do not duplicate results described elsewhere in the article.

Figures. Diagrams, charts, line drawings, chemical formulas and photographs should be referred to as figures. Figures should be drawn professionally. Do not put the title of the figure in the figure! Each axis of charts should be labeled with a description of variable it represents. Only the first letter of the first word should be capitalized, and size for letters should be 10 points. Charts should be in gray scale.

Photographs, chemical formulas and diagrams should be saved as separate, stand-alone files and not embedded as part(s) of the text file. The files containing photographs, chemical formulas, etc. should be named with the number of the figure in the text (e.g. Figure 5). It is preferable to submit images, created at 100% image size. File formats in order of preference are JPEG, TIFF, BMP; always use a minimum of 300 dpi. The title and the legend of figures should be given in the text file after tables and graphs.

Optional suggestions for the referees (excluding co-authors or collaborators for the last three years) of the paper, with the complete mailing address, e-mail address, phone and fax numbers.

Manuscripts are not returned to authors.

All correspondence, including notification of the Editor's decision and requests for revisions, will be sent to you by e-mail.

ETHICAL GUIDELINES

Based on Uniform Requirements for Manuscripts Submitted to Biomedical Journals; International Committee of Medical Journal Editors www.ICMJE.org

Authorship

An author is a physical person, whose creative activity has resulted in a work. The authorship of a work, created by two or more persons, is owned by them, irrespective of the fact that the work constitutes one whole or is comprised of independent parts, and the creators are considered co-authors.

Authorship constitutes the basis for evaluation of responsibility and credit for intellectual work. Authorship is important for the reputation, academic promotion, and granting support to the individuals involved, as well as for the strength and reputation of their institution. Authorship practices should reflect actual contribution to the creation of a final product.

Authorship credit is granted if the author has made a considerable contribution to conception and design, data collection, or analysis and interpretation of information, important contents has been found after critical revision and it has been finally approved for publication.

The final version of the manuscript should be approved by all co-authors, and their consent to submission for publication is to be stated by completing the SUBMISSION FORM.

All contributors who do not meet authorship criteria should be mentioned in an acknowledgments section, for example persons who have provided technical or writing assistance by collecting data, or a unit's chairperson who has solely been generally supportive. Acknowledgment(s) for financial and material support should also be included.

Editorship

The editor-in-chief of Journal of Biomedical & Clinical Research has full authority over the contents of the journal and the timing of publication of that contents. The concept of editorial freedom should be resolutely defended by editors.

An independent editorial advisory board helps the editor establish and maintain editorial policy. The validation of the work in question and its importance to researchers and readers must always drive the editor's decisions. The editor is guided by the policies of the journal's editorial board and constrained by such legal requirements as shall then be in force regarding libel, copyright infringement and plagiarism. The editor may confer with the members of editorial board or reviewers in making this decision.

Peer Review

Journal of Biomedical & Clinical Research aims at publishing quality research while maintaining rigorous peer review process. Manuscripts that meet quality and technical requirements of the journal will be peer-reviewed by two or more experts in the relevant fields, and a statement will be sent authors in about two months. They will be notified by e-mail, if the review process take more time. Manuscripts with noteworthy results will be reviewed and published at the highest priority. Possible decisions regarding a manuscript are acceptance, acceptance after minor revision, acceptance after major revision or rejection. If minor revision is necessary, authors should send back their revised version as soon as possible but not later than 15 days. If major revision is required, authors are expected to return a corrected manuscript within 30 days.

Conflicts of Interest

All authors should disclose any financial or other significant conflict of interest that might affect the results or interpretation of their manuscript. They can provide additional details, if necessary, in a separate attachment, accompanying the manuscript All sources of financial support for the project should be disclosed. Potential conflicts of interest which should be disclosed include employment, consultancies, stock ownership, honoraria, patent applications/registrations, and grants or other funding. Potential conflicts of interest should be disclosed at the earliest stage possible.

Editors are expected to avoid selecting peer reviewers should there be doubts about potential conflicts of interest. Reviewers should inform the editors about any conflicts of interest that could bias their evaluation of the manuscript, and they should refuse to review a manuscripts in case of potential bias. Editors who finally decide about manuscripts are to avoid personal, professional, or financial implications in any of the issues they might arbitrate.

Privacy and Confidentiality

Authors' confidentiality should be protected during the process of reviewing a manuscript. By submitting a manuscripts for review, authors entrust editors with results of their scientific research and creative effort. Breaking confidentiality while reviewing is in process may violate the author's rights and might be unfavourable to their reputation and career. Reviewers' confidentiality should be also respected by the editor. Confidentiality must be honored except in cases of alleged dishonesty or fraud.

Editors should not share information about manuscripts regarding receipt, content, status in the reviewing process, criticism by reviewers, or final decisions on publishing, with anyone other than the corresponding author, reviewers, potential reviewers, other editorial advisers, and the publisher, as appropriate, including requests for using the materials in legal proceedings.

Reviewers should also be notified of the editor's decision on acceptance or rejection of a manuscript.

Protection of Human Subjects and Animals in Research

When reporting experiments on humans, authors are expected to indicate whether the procedures followed were in accordance with ethical standards of institutional and national committees responsible for human experimentation and in accordance with the Helsinki Declaration of 1975, as revised in 2008. If doubt exists whether the research was carried out in accordance with the Helsinki Declaration, authors should provide explanations about the rationale for their approach and demonstrate explicit approval by the institutional review body of doubtful aspects of the study. In reports on experiments on animals, authors should indicate compliance with institutional and national guide for care and use of laboratory animals.

Duplicate publication

Manuscripts submitted for publication in Journal of Biomedical & Clinical Research must not have been published or submitted for publication elsewhere. Authors who have submitted manuscripts for consideration or in press elsewhere should explicitly present any relevant current information in a cover letter to the attention of the editor. If part of a contribution submitted to Journal of Biomedical & Clinical Research has appeared or is expected to appear elsewhere, the details must be specified by the author in a cover letter.

Authors are responsible for obtaining permission from a publisher or coryright holder for reproduction of figures or tables previously published elsewhere .

The editors consider all material in good faith that the journal has full permission to publish every part of the submitted material, including illustrations.

Plagiarism

Plagiarism implies using language and ideas of another author or their close imitation and presenting them as one's own original work. Duplicate publication, sometimes called self-plagiarism, occurs when an author uses substantial parts of his or her own published work without appropriate referencing. This includes submitting a paper published in other journals, to which authors have added small amounts of new data or a previously published paper.

Plagiarism is considered evident in case when large chunks of text have been cut and pasted. Such manuscripts would not be considered for publication in Journal of Biomedical & Clinical Research. However, examples of minor plagiarism without dishonest intent are relatively frequent, as us the case of reusing parts of materials and methods or an introduction from an earlier paper. Any such will be judged by the editors on its own merits.

In case plagiarism is found, the journal will contact the author's institute. The paper containing the plagiarism will also be obviously marked or, depending on the extent of the plagiarism, the paper may also be formally withdrawn.

SUBMISSION FORM

TITLE:

A signature below Certifies Compliance with the Following Statements

Copyright Transfer. In consideration of the acceptance of the above work for publication, I do hereby assign and transfer to Journal of Biomedical and Clinical Research all rights, title, and interest in and to the copyright in the above-titled work. This includes preliminary display/posting of the abstract of the accepted article in electronic form before publication.

Authorship Responsibilities. I attest:

- 1. The manuscript is not published previously and under consideration for publication elsewhere.
- 2. The manuscript is truthful original work/review without fabrication or plagiarism.
- 3. I have made an important scientific contribution to the study and I am absolutely familiar with the primary data.
- 4. I have read the complete manuscript and take responsibility for the content and completeness if the manuscript.

Conflict of Interest Disclosure. All funding sources supporting the work and all institutional affiliations of mine are acknowledged. Except as disclosed on a separate attachment, I certify that I have no commercial associations (e.g. consultancies, ownership, equity interests, patient-licensing arrangements) that might pose a conflict of interest in connection with the submitted article (letter attached).

Signature from	author	required	in t	the	same	order	as	on	the	manusc	ript	title
page												

Signature (1)	Name	Date
Signature (2)	Name	Date
Signature (3)	Name	Date
Signature (4)	Name	Date
Signature (5)	Name	Date
Signature (6)	Name	Date
Signature (7)	Name	Date
Signature (8)	Name	Date
Signature (9)	Name	Date

Corresponding Author

Phone

Address e-mail